


ForgetMeNot - ref.: FMNY2B (yellow) or FMNB2B (blue)

Simple answer to a real issue*: digit tourniquet ForgetMeNot

Available in yellow or blue Sold in packs of 10

Reusable Sold no sterile





* Résultats fonctionnels à long terme d'ischémies digitales sous garrot: à propos de 2 observations. SOFCOT 2010 / Florence Mallard (Angers) Julien Richou, Dominique Le Nen



Quick video to present exsanguination

by Stéphanie Gouzou - Strasbourg University Hospital

CLICK HERE

http://www.arex.fr/QRc.php?c=23

 $C \in$ AREX® 3 allée du Clos Tonnerre 91120 Palaiseau - FRANCE

www.arex.fr

ForgetMeNot

The only specifically designed-for-purpose, reusable finger tourniquet.

Reusable

One size-fits-all

Achieving a totally exsanguinated finger

Bright blue or yellow highly visible tails make it totally unforgettable

Removable without rolling it over the wound or any risk of cutting skin

Environmentally friendly since it is reusable

For optimal FMN use, follow the detailed instructions below. Placement of the FMN tourniquet is simple, easy and repeatable. It is used under local (LA) or locoregional (LRA) anaesthesia.

1. The tourniquet is prepared before placement by forming one or two loops, passing one, or both (as preferred), silicone beaded strand(s) through holes in the central crosspiece



2. The tourniquet is then placed on the digital pulp and tightened by pulling on the two strands.





6. Removing the FMN is easy: simply pull on the central crosspiece to loosen and remove the tourniquet. The FMN can be sterilised

3. At this point, the tourniquet is

guination efficacy.

ness during surgery.

rolled down towards the base

of the finger to obtain exsan-

If during this maneuvre the FMN slip over skin without wrapping around itself then tourniquet tightness may not be maintained. FMN has to wrap around itself to reach finger base and then to maintain finger vascular tight-



When a totally bloodless field is not required, the FMN can be used with a single loop: the arm is held above heart level for a few minutes and the FMN is applied directly to the base of the finger.

- NB1 Do not use ForgetMeNot on small fingers.
- NB2 Do not use ForgetMeNot in paediatric cases.

These contra-indication are related to FMN dimentional characteristics.

NB3 - Do not clamp ForgetMeNot with a metallic instrument to maintain this finger tourniquet, this may initiate small cuts into silicone resulting in potential secondary rupture.